



TNC priority areas
 Marten buffer
 NECEC

Marten habitat suitability

High
 Moderate
 Intermediate
 Low
 Areas of broad-scale connectivity

Overview map showing the total area encompassed in the analysis and the 4 map subsets. In addition, areas with higher broad-scale connectivity of marten habitat are identified by 3 black hatched polygons. Areas with High habitat suitability for American Marten were mature, intact forest ca. 2018. Paritally-harvested forest is considered Moderate suitability. Areas with Intermediate suitability are older (1970s-1980s) regenerating clearcuts. Areas with Low suitability are younger (1990s-2018) regenerating clearcuts. Marten buffer width is a total of 6000 ft, corresponding to the width of an average home range size for a female marten in Maine (650 ac), assuming a circular home range.

Prepared for DEP by E. Simons-Legaard (5/17/2019)





TNC priority areas
 Marten buffer
 NECEC

Marten habitat suitability
 High
 Moderate
 Intermediate
 Low

Areas with High habitat suitability for American Marten were mature, intact forest ca. 2018. Paritally-harvested forest is considered Moderate suitability. Areas with Intermediate suitability are older (1970s-1980s) regenerating clearcuts. Areas with Low suitability are younger (1990s-2018) regenerating clearcuts. Marten buffer width is a total of 6000 ft, corresponding to the width of an average home range size for a female marten in Maine (650 ac), assuming a circular home range.

Prepared for DEP by E. Simons-Legaard (5/17/2019)




TNC priority areas
 Marten buffer
 NECEC

Marten habitat suitability

High
 Moderate
 Intermediate
 Low

Areas with High habitat suitability for American Marten were mature, intact forest ca. 2018. Paritally-harvested forest is considered Moderate suitability. Areas with Intermediate suitability are older (1970s-1980s) regenerating clearcuts. Areas with Low suitability are younger (1990s-2018) regenerating clearcuts. Marten buffer width is a total of 6000 ft, corresponding to the width of an average home range size for a female marten in Maine (650 ac), assuming a circular home range.

Prepared for DEP by E. Simons-Legaard (5/17/2019)





TNC priority areas
 Marten buffer
 NECEC

Marten habitat suitability

High
 Moderate
 Intermediate
 Low

Areas with High habitat suitability for American Marten were mature, intact forest ca. 2018. Paritally-harvested forest is considered Moderate suitability. Areas with Intermediate suitability are older (1970s-1980s) regenerating clearcuts. Areas with Low suitability are younger (1990s-2018) regenerating clearcuts. Marten buffer width is a total of 6000 ft, corresponding to the width of an average home range size for a female marten in Maine (650 ac), assuming a circular home range.



TNC priority areas
 Marten buffer
 NECEC

Marten habitat suitability

High
 Moderate
 Intermediate
 Low

Areas with High habitat suitability for American Marten were mature, intact forest ca. 2018. Paritally-harvested forest is considered Moderate suitability. Areas with Intermediate suitability are older (1970s-1980s) regenerating clearcuts. Areas with Low suitability are younger (1990s-2018) regenerating clearcuts. Marten buffer width is a total of 6000 ft, corresponding to the width of an average home range size for a female marten in Maine (650 ac), assuming a circular home range.

Prepared for DEP by E. Simons-Legaard (5/17/2019)