


# CLIMATE CHANGE PROJECTIONS FOR INDIVIDUAL TREE SPECIES EASTERN AND COASTAL MAINE



This region's forests will be affected by a changing climate and other stressors during this century. A team of managers and researchers created an assessment that describes the vulnerability of forests in the region ([Janowiak et al. 2018](#)). This report includes information on observed and future climate trends, and also summarizes key vulnerabilities

for forested natural communities. The Landscape Change Research Group recently updated the Climate Change Tree Atlas, and this handout summarizes that information. Full Tree Atlas results are available online at [www.fs.fed.us/nrs/atlas/](http://www.fs.fed.us/nrs/atlas/). Two climate scenarios are presented to "bracket" a range of possible futures. These future climate projections (2070 to 2099) provide information about how individual tree species may respond to a changing climate. Results for "low" and "high" emissions scenarios can be compared on the reverse side of this handout.

The updated Tree Atlas presents additional information helpful to interpret tree species changes:

- **Suitable habitat** - calculated based on 39 variables that explain where optimum conditions exist for a species, including soils, landforms, and climate variables.
- **Adaptability** - based on life-history traits that might increase or decrease tolerance of expected changes, such as the ability to withstand different forms of disturbance.
- **Capability** - a rating of the species' ability to cope or persist with climate change in this region based on suitable habitat change (statistical modeling), adaptability (literature review and expert opinion), and abundance (FIA data). The capability rating is modified by abundance information; ratings are downgraded for rare species and upgraded for abundant species.
- **Migration Potential Model** - when combined with habitat suitability, an estimate of a species' colonization likelihood for new habitats. This rating can be helpful for assisted migration or focused management (see the table section: "New Habitat with Migration Potential").

Remember that models are just tools, and they're not perfect. Model projections can't account for all factors that influence future species success. If a species is rare or confined to a small area, model results may be less reliable. These factors, and others, could cause a particular species to perform better or worse than a model projects. Human choices will also continue to influence forest distribution, especially for tree species that are projected to increase. Planting programs may assist the movement of future-adapted species, but this will depend on management decisions. Despite these limits, models provide useful information about future expectations. It's perhaps best to think of these projections as indicators of possibility and potential change.

**SOURCE:** This handout summarizes the full model results for Eastern and Coastal Maine, available at [www.fs.fed.us/nrs/atlas/combined/resources/summaries](http://www.fs.fed.us/nrs/atlas/combined/resources/summaries). More information on vulnerability and adaptation in the New England region can be found at [www.forestadaptation.org/new-england](http://www.forestadaptation.org/new-england). A full description of the models and variables are provided in Iverson et al. 2019 ([www.nrs.fs.fed.us/pubs/57857](http://www.nrs.fs.fed.us/pubs/57857) and [www.nrs.fs.fed.us/pubs/59105](http://www.nrs.fs.fed.us/pubs/59105)) and Peters et al. 2019 ([www.nrs.fs.fed.us/pubs/58353](http://www.nrs.fs.fed.us/pubs/58353)).

## CLIMATE CHANGE CAPABILITY

### POOR CAPABILITY

American mountain-ash	Pitch pine
Balsam poplar	Red pine
Black spruce	Serviceberry
Bur oak	Striped maple
Jack pine	Tamarack (native)
Mountain maple	White spruce
Pin cherry	

### FAIR CAPABILITY

Balsam fir	Northern white-cedar
Eastern hemlock	Red spruce
Gray birch	

### GOOD CAPABILITY

American beech	Northern red oak
American elm	Quaking aspen
Bigtooth aspen	Red maple
Black cherry	Sugar maple
Black oak	Swamp white oak
Ironwood	White oak

### MIXED RESULTS

American basswood	Green ash
American hornbeam	Paper birch
Black ash	Silver maple
Boxelder	White ash
Eastern white pine	Yellow birch

### NEW HABITAT WITH MIGRATION POTENTIAL

Bitternut hickory	Pignut hickory
Black walnut	Sassafras
Blackgum	Scarlet oak
Chestnut oak	Shagbark hickory
Eastern redcedar	Slippery elm
Mockernut hickory	Yellow-poplar



**ADAPTABILITY:** Life-history factors, such as the ability to respond favorably to disturbance, that are not included in the Tree Atlas model and may make a species more or less able to adapt to future stressors.

- + **HIGH** Species may perform better than modeled
- **MEDIUM**
- **LOW** Species may perform worse than modeled

**HABITAT CHANGE:** Projected change in suitable habitat between current and potential future conditions.

- ▲ **INCREASE** Projected increase of >20% by 2100
- **NO CHANGE** Projected change of <20% by 2100
- ▼ **DECREASE** Projected decrease of >20% by 2100
- ★ **NEW HABITAT** Tree Atlas projects new habitat for species not currently present

**ABUNDANCE:** Based on Forest Inventory Analysis (FIA) summed Importance Value data, calibrated to a standard geographic area.

- + **ABUNDANT**
- **COMMON**
- **RARE**

**CAPABILITY:** An overall rating that describes a species' ability to cope or persist with climate change based on suitable habitat change class (statistical modeling), adaptability (literature review and expert opinion), and abundance within this region.

- ▲ **GOOD** Increasing suitable habitat, medium or high adaptability, and common or abundant
- **FAIR** Mixed combinations, such as a rare species with increasing suitable habitat and medium adaptability
- ▼ **POOR** Decreasing suitable habitat, medium or low adaptability, and uncommon or rare

SPECIES	LOW CLIMATE CHANGE (RCP 4.5)				HIGH CLIMATE CHANGE (RCP 8.5)		SPECIES	LOW CLIMATE CHANGE (RCP 4.5)				HIGH CLIMATE CHANGE (RCP 8.5)	
	ADAPT	ABUN	HABITAT CHANGE	CAPABILITY	HABITAT CHANGE	CAPABILITY		ADAPT	ABUN	HABITAT CHANGE	CAPABILITY	HABITAT CHANGE	CAPABILITY
American basswood	•	-	●	▼	▲	▲	Northern red oak	+	•	▲	▲	▲	▲
American beech	•	+	●	▲	●	▲	Northern white-cedar	•	+	▼	○	▼	○
American elm	•	-	▲	▲	▲	▲	Overcup oak	-	-	★		★	
American hornbeam*	•	-	●	▼	▲	▲	Paper birch	•	•	●	○	▼	▼
American mountain-ash*	-	-	▼	▼	▼	▼	Pignut hickory	•	-	★		★	
Balsam fir	-	+	▼	○	▼	○	Pin cherry*	•	-	▼	▼	▼	▼
Balsam poplar	•	•	▼	▼	▼	▼	Pin oak*	-	-	★		★	
Bigtooth aspen	•	•	▲	▲	▲	▲	Pitch pine	•	-	●	▼	●	▼
Bitternut hickory*	+	-	★		★		Post oak	+	-	★		★	
Black ash	-	•	▲	○	●	▼	Quaking aspen	•	•	▲	▲	▲	▲
Black cherry	-	•	▲	▲	▲	▲	Red maple	+	+	▲	▲	●	▲
Black locust*	•	-	★		★		Red pine	-	-	●	▼	●	▼
Black oak	•	-	▲	▲	▲	▲	Red spruce	-	+	▼	○	▼	○
Black spruce	•	•	▼	▼	▼	▼	Sassafras*	•	-	★		★	
Black walnut*	•	-	★		★		Scarlet oak	•	-	★		★	
Blackgum	+	-	★		★		Serviceberry*	•	-	▼	▼	●	▼
Boxelder*	+	-	●	○	▲	▲	Shagbark hickory	•	-	★		★	
Bur oak	+	-	▼	▼	▼	▼	Silver maple*	+	-	▼	▼	▲	▲
Chestnut oak	+	-	★		★		Slippery elm*	•	-			★	
Common persimmon*	+	-			★		Striped maple	•	•	▼	▼	▼	▼
Cucumbertree*	•	-			★		Sugar maple	+	•	▲	▲	▲	▲
Eastern hemlock	-	+	●	○	▼	○	Swamp white oak*	•	-	▲	▲	▲	▲
Eastern redcedar	•	-	★		★		Sweet birch	-	-	★		★	
Eastern white pine	-	+	▲	▲	●	○	Sweetgum	•	-	★		★	
Flowering dogwood	•	-			★		Sycamore*	•	-	★		★	
Gray birch*	•	•	●	○	●	○	Tamarack (native)	-	•	▼	▼	▼	▼
Green ash*	•	-	●	▼	▲	▲	Virginia pine	•	-	★		★	
Hackberry	+	-			★		White ash	-	•	▲	○	▲	▲
Ironwood*	+	•	▲	▲	▲	▲	White oak	+	-	▲	▲	▲	▲
Jack pine	+	-	▼	▼	▼	▼	White spruce	•	•	▼	▼	▼	▼
Mockernut hickory	+	-	★		★		Yellow birch	•	•	●	○	▼	▼
Mountain maple*	+	-	▼	▼	▼	▼	Yellow-poplar	+	-	★		★	

\*Species with low model reliability based on five statistical metrics of the habitat models that affect change class. See maps and tables for more information ([www.fs.fed.us/nrs/atlas/combined/resources/summaries](http://www.fs.fed.us/nrs/atlas/combined/resources/summaries)).