

# Is it safe for my child to get these vaccines at the same time?



	COVID-19 (RNA)	Influenza (Flu)
Diphtheria, tetanus, pertussis (DTaP) vaccine	✓	*
<i>Haemophilus influenzae</i> B vaccine (Hib)	✓	✓
Hepatitis A vaccine	✓	✓
Hepatitis B vaccine	✓	✓
Human papillomavirus vaccine (HPV)	✓	✓
Measles, mumps, rubella vaccine (MMR)	✓	✓
Meningococcal serogroup ACWY vaccine (MenACWY)	✓	✓
Meningococcal serogroup B vaccine (MenB)	✓	✓
Pneumococcal vaccine(s)	✓	*
Polio vaccine	✓	✓
Rotavirus vaccine	✓	✓
Tetanus, diphtheria, pertussis (Tdap) vaccine	✓	✓
Varicella vaccine	✓	✓

\* Talk to a healthcare provider about getting these vaccines at the same time as the influenza vaccine. A federal CDC study showed a small increased risk for febrile seizures during the 24 hours after a child receives the inactivated influenza vaccine at the same time as the PCV13 vaccine or DTaP vaccine. However, the risk of febrile seizure with any combination of these vaccines is small and ACIP recommends giving these vaccines at the same visit if indicated. See [www.cdc.gov/vaccinesafety/concerns/febrile-seizures.html](http://www.cdc.gov/vaccinesafety/concerns/febrile-seizures.html) or call Maine CDC at 1-800-821-5821 for more information.